ScyTek Laboratories	Instructions For Use TM3-IFU		
	Rev. Date: 10/29/03	Revision: 2	Page 1 of 1

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Fax (435) 755-0015 - www.scytek.com

TMB Precipitating Reagent (Standard Sensitivity)

Description: This liquid substrate for peroxidase consists of tetramethylbenzidine (TMB) plus dilute hydrogen peroxide in a single-reagent stabilized form. The reagent has been specifically formulated for measuring peroxidase in EIA systems. This reagent is stable for long-term storage, and provides sensitivity equal to, or greater than, that of other commercially available chromogens.

Form: 3,3',5,5'-tetramethylbenzidine

Contents: TMB in a dilute organic solvent, pH 6.0+ 0.5. Hydrogen peroxide is added at a concentration of 0.03%

- **Storage:** Store TMB solution at 2 8°C. Storage below this temperature may result in a slight blue tint, which disappears after returning to the appropriate temperature. Refrigeration is not mandatory. Do not expose reagents to strong light. Keep bottle tightly capped. Reagents are stable for a least 12 months when stored at room temperature, or 18 months when stored at 2 8°C. Avoid contamination of reagents with labware which has not been thoroughly cleaned. A slight yellow tinge may develop over time. This does not affect product performance. Do not use if solution darkens.
- Activating Agents: Peroxidase
- Light Sensitivity: Negligible for short exposure times
- Reaction Time: Approximately 15 minutes (Range 5 60 min.)
- Reaction pH: Approximately pH 6.0 (Range 5.0 7.0)
- Reaction Temperature: Room temperature
- **Precautions:** Do not pipet reagents by mouth. Avoid contact with skin and eyes. In case of contact, flush with plenty water. Dispose of reagents according to local regulations. If disposed of in the sink, flush with plenty of water.