

## Instructions For Use A00073-IFU

Rev. Date: 10/15/03

**Revision: 2** 

Page 1 of 1

P.O. Box 3286 - Logan, Utah 84323, U.S.A. - Tel. (800) 729-8350 - Fax (435) 755-0015 - www.scytek.com

## Polyclonal Rabbit Antibody to Human Alpha-1-Antichymotrypsin

## **Description:**

Immunogen: Alpha-1-Antichymotrypsin from human serum.

Format: This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-

embedded tissue sections. No further titration is required.

Specificity: This antibody recognizes macrophages and monocytes in human colon, tonsil and skin.

Species Reactivity: Human. Shows no cross-reactivity with horse, sheep, swine, goat, cow, dog, cat,

chicken, rat, mouse, mink, kangaroo or Guinea pig. Others-not tested.

Positive Control: Tonsil.

Uses/Limitations: Immunohistochemistry

For Research Use Only.

Do not use past expiration date.

**Storage:** 2-8° Centigrade.

**Procedure:** ScyTek suggests an incubation period of 30 minutes at room temperature. However,

depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>Enzymatic predigestion of paraffin</u>

embedded tissues may improve immunostaining, but is not required.

**Precautions:** Contains Sodium Azide as a preservative.

Do not pipette by mouth.

**References:** i) Meis JM, Osborne BM, Butler JJ. A comparative marker study of large cell lymphoma,

Hodgkin's disease, and true histiocytic lymphoma in paraffin-embedded tissue. American

Journal of Clinical Pathology 1986; 86:591-599.

ii) Nathrath WBJ, Meister P. Lysozyme (muramidase) and alpha-1-antichymotrypsin as immunohistochemical tumour markers. Acta Histochem 1982; (suppl 25): 69-72.