


# CD106 / VCAM1 (Activated Endothelial Cell Marker); Clone VCAM1/843 (Concentrate)


| <b>Availability/Contents:</b> | <u>Item #</u>   | <u>Volume</u>          |
|-------------------------------|---|------------------------|
|                               | RA0336-C.5  | 0.5 ml                 |
| <b>Description:</b>           |   |                        |
| Species:                      | Mouse   |                        |
| Immunogen:                    | Recombinant human VCAM1 protein   |                        |
| Clone:                        | VCAM1/843   |                        |
| Isotype:                      | IgG1, kappa   |                        |
| Entrez Gene ID:               | 7412 (Human)  |                        |
| Hu Chromosome Loc.:           | 1q21.2  |                        |
| Synonyms:                     | CD106; INCAM-100; L1CAM; Vascular Cell Adhesion Molecule 1 (VCAM1); Vascular cell adhesion protein 1  |                        |
| Mol. Weight of Antigen:       | 110kDa  |                        |
| Format:                       | 200µg/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.  |                        |
| Specificity:                  | Recognizes a protein of 110kDa, identified as CD106 (also known as vascular cell adhesion molecule-1 (VCAM-1) and INCAM-100).   |                        |
| Background:                   | CD106 is a member of the Ig superfamily of adhesion molecules and is expressed at high levels on cytokine stimulated vascular endothelial cells, and at minimal levels on un-stimulated endothelial cells. It is also present on follicular and inter-follicular dendritic cells of lymph nodes, myoblasts, and some macrophages. CD106 serves as a ligand for leukocyte integrin alpha4beta1 (VLA-4 or CD49d/CD29) and mediates cell adhesion of leukocytes to activated endothelium. It plays a role in various immunological and inflammatory responses. |                        |
| Species Reactivity:           | Human. Others not known.  |                        |
| Positive Control:             | Human placenta or tonsil.   |                        |
| Cellular Localization:        | Cell surface  |                        |
| Titer/ Working Dilution:      | Immunohistochemistry (Frozen and Formalin-fixed): 0.5-1 µg/ml   |                        |
|                               | Flow Cytometry:   | 0.5-1 µg/million cells |
|                               | Immunofluorescence:   | 1-2 µg/ml              |
| Microbiological State:        | This product is not sterile.  |                        |

Storage: 2° C  8° C




ScyTek Laboratories, Inc.  
 205 South 600 West  
 Logan, UT 84321  
 U.S.A.



 EmergoEurope (31)(0) 70 345-8570  
 Molsnstraat 15  
 2513 BH Hague, The Netherlands

**Uses/Limitations:** Not to be taken internally.  
 For Research Use Only.  
 This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.  
 Do not use if reagent becomes cloudy.  
 Do not use past expiration date.  
 Non-Sterile.



Formalin-fixed, paraffin-embedded human tonsil stained with CD106; Clone VCAM1/843.

**Ordering Information and Current Pricing at [www.scytek.com](http://www.scytek.com)**

**Procedure:**


1. **Tissue Section Pretreatment (Required):** Staining of formalin fixed, paraffin embedded tissue sections is significantly enhanced by pretreatment with EDTA Buffer (10X) HIER Solution (pH 8.0) (ScyTek catalog# ETA).
2. **Primary Antibody Incubation Time:** We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user.
3. **Visualization:** For maximum staining intensity we recommend the “UltraTek HRP Anti-Polyvalent Lab Pack” (ScyTek catalog# UHP125, see IFU for instructions) combined with the “DAB Chromogen/Substrate Bulk Pack (High Contrast)” (ScyTek catalog# ACV500, see IFU for instructions).


**Precautions:** Contains Sodium Azide as a preservative (0.09% w/v).  
 Do not pipette by mouth.  
 Avoid contact of reagents and specimens with skin and mucous membranes.  
 Avoid microbial contamination of reagents or increased nonspecific staining may occur.  
 This product contains no hazardous material at a reportable concentration according to U.S. 29 CFR 1910.1200, OSHA Hazardous Communication Standard and EC Directive 91/155/EC.

**References:**

1. Vermot-Desroches C et al. Heterogeneity of antigen expression among human umbilical cord vascular endothelial cells: identification of cell subsets by co-expression of haemopoietic antigens. *Immunol Lett* 1995, 48(1):1-9
2. Rice GE et al. Vascular and nonvascular expression of INCAM-110. A target for mononuclear leukocyte adhesion in normal and inflamed human tissues. *Am J Pathol* 1991, 138(2):385-393.
3. Huang MJ et al. Expression of vascular cell adhesion molecule-1 by follicular dendritic cells. *Leuk Lymphoma* 1995, 18(3-4):259-264.


**Warranty:** No products or “Instructions For Use (IFU)” are to be construed as a recommendation for use in violation of any patents. We make no representations, warranties or assurances as to the accuracy or completeness of information provided on our IFU or website. Our warranty is limited to the actual price paid for the product. ScyTek Laboratories, Inc. is not liable for any property damage, personal injury, time or effort or economic loss caused by our products. Immunohistochemistry is a complex technique involving both histological and immunological detection methods. Tissue processing and handling prior to immunostaining can cause inconsistent results. Variations in fixation and embedding or the inherent nature of the tissue specimen may cause variations in results. Endogenous peroxidase activity or pseudoperoxidase activity in erythrocytes and endogenous biotin may cause non-specific staining depending on detection system used.

Storage: 2° C  8° C



ScyTek Laboratories, Inc.  
 205 South 600 West  
 Logan, UT 84321  
 U.S.A.



 EmergoEurope (31)(0) 70 345-8570  
 Molsnstraat 15  
 2513 BH Hague, The Netherlands